

TexaClear® Allergy Relief Drug Facts

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product package labeling, this website labeling should be regarded as the most current.

Drug Facts	
Active ingredients (in each 30 mL (2 TBSP))	Purpose
Chlophedianol HCl 25 mg.....	Cough suppressant
Pyrilamine Maleate 50 mg	Antihistamine
Uses	
temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:	
<ul style="list-style-type: none"> ▪ cough due to minor throat and bronchial irritation ▪ runny nose ▪ sneezing ▪ itching of the nose or throat ▪ itchy, watery eyes 	
Warnings	
Do not exceed recommended dosage.	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> ▪ a cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema ▪ a cough that occurs with too much phlegm (mucus) ▪ glaucoma ▪ a breathing problem such as emphysema or chronic bronchitis ▪ difficulty in urination due to enlargement of the prostate gland 	
Ask a doctor or pharmacist before use if you are ▪ taking sedatives or tranquilizers.	
When using this product	
<ul style="list-style-type: none"> ▪ excitability may occur, especially in children ▪ marked drowsiness may occur ▪ avoid alcoholic drinks ▪ alcohol, sedatives, and tranquilizers may increase drowsiness ▪ be careful when driving a motor vehicle or operating machinery 	
Stop use and ask a doctor if:	
<ul style="list-style-type: none"> ▪ nervousness, dizziness or sleeplessness occurs ▪ symptoms do not improve within 7 days, tends to recur, or are accompanied by a fever, rash or persistent headache. A persistent cough may be a sign of a serious condition. ▪ new symptoms occur 	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions	
<ul style="list-style-type: none"> ▪ do not exceed recommended dosage. ▪ do not take more than 4 doses in any 24-hour period. ▪ use enclosed dose cup ▪ keep dosage cup with product ▪ TBSP= tablespoon ▪ mL= milliliter 	
adults and children 12 years of age and over:	30 mL (2 TBSP) every 6-8 hours.
children under 12 years of age:	do not use.
Other information	
<ul style="list-style-type: none"> ▪ each 30 mL (2 TBSP) contains: Sodium 13 mg. ▪ tamper evident: Do not use if foil seal under cap is broken or missing. ▪ read all product information before using ▪ store at 68° to 86°F (20° to 30°C). ▪ avoid excessive heat and humidity 	
Inactive ingredients	
anhydrous citric acid, glycerin, propylene glycol, purified water, sodium citrate, sodium saccharin, sorbitol	
Questions? Comments?	
Call 1-888-535-0305 9 a.m. - 5 p.m. CST.	



Distributed by:
GM Pharmaceuticals, Inc.
Arlington, TX. 76015

TexaClear® Allergy Relief One Dose Drug Facts

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product package labeling, this website labeling should be regarded as the most current.

Drug Facts	
Active ingredients (per 57 mL)	Purpose
Chlophedianol HCl 25 mg.....	Cough suppressant
Pyrilamine Maleate 50 mg	Antihistamine
Uses	
temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:	
<ul style="list-style-type: none"> ▪ cough due to minor throat and bronchial irritation ▪ runny nose ▪ sneezing ▪ itching of the nose or throat ▪ itchy, watery eyes 	
Warnings	
Do not exceed recommended dosage.	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> ▪ a cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema ▪ a cough that occurs with too much phlegm (mucus) ▪ glaucoma ▪ a breathing problem such as emphysema or chronic bronchitis ▪ difficulty in urination due to enlargement of the prostate gland 	
Ask a doctor or pharmacist before use if you are - taking sedatives or tranquilizers.	
When using this product	
<ul style="list-style-type: none"> ▪ do not use more than directed ▪ excitability may occur, especially in children ▪ marked drowsiness may occur ▪ avoid alcoholic drinks ▪ alcohol, sedatives, and tranquilizers may increase drowsiness ▪ be careful when driving a motor vehicle or operating machinery 	
Stop use and ask a doctor if:	
<ul style="list-style-type: none"> ▪ nervousness, dizziness or sleeplessness occurs ▪ symptoms do not improve within 7 days, tends to recur, or are accompanied by a fever, rash or persistent headache. A persistent cough may be a sign of a serious condition. ▪ new symptoms occur 	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions	
<ul style="list-style-type: none"> ▪ do not use more than directed ▪ do not use in children under 12 years of age ▪ adults and children 12 years of age and over: take 1 (57 mL) bottle every 6 to 8 hours. ▪ do not take more than 4 (57 mL) bottles in 24 hours unless directed by a doctor 	
Other information	
<ul style="list-style-type: none"> ▪ each 57 mL bottle contains: Sodium 24 mg ▪ tamper evident: Do not use if neckband on bottle cap is broken or missing. ▪ read all product information before using ▪ store at 68° to 86°F (20° to 30°C). ▪ avoid excessive heat and humidity 	
Inactive ingredients	
citric acid anhydrous, glycerin, propylene glycol, purified water, sodium citrate, sodium saccharin, sorbitol	
Questions? Comments?	
Call 1-888-535-0305 9 a.m. - 5 p.m. CST.	



Distributed by:
GM Pharmaceuticals, Inc.
Arlington, TX. 76015